

Hamburger Beiträge

zur Angewandten Mathematik

Linear equations and the Kronecker product in coquaternions

Drahoslava Janovská and Gerhard Opfer

Dedicated to Lothar Reichel on the occasion of his 60th birthday

Dieser Artikel wird – in modifizierter Form – 2013 unter dem angegebenen Titel
erscheinen in den Mitteilungen der Mathematischen Gesellschaft in Hamburg
(Mitt. Math. Ges. Hamburg).

Nr. 2013-05
May 2013

LINEAR EQUATIONS AND THE KRONECKER PRODUCT IN COQUATERNIONS

DRAHOSLAVA JANOVSKÁ* AND GERHARD OPFER†

Dedicated to Lothar Reichel on the occasion of his 60th birthday

Abstract. We present the solution of the general linear system in n coquaternionic variables and m equations, and derive the Kronecker product from the matrix system \mathbf{AXB} in coquaternions. It is shown, that the coquaternionic Kronecker product can be computed by means of dyadic coquaternionic products. The special matrix case $\mathbf{Ax} = \mathbf{b}$ is also included. The one dimensional case, including Sylvester's equation is solved and there are several, nontrivial, numerical examples.

Key words: Coquaternions, split quaternions, linear systems of equations in coquaternions, $\mathbf{Ax} = \mathbf{b}$ in coquaternions, Kronecker product in coquaternions, dyadic coquaternionic products, Sylvester's equation in coquaternions.

AMS Subject classification: 11R52, 12E15, 12Y05, 15A06, 15A24.

1. Introduction. Coquaternions were introduced, 1849, by Sir James Cockle (1819–1895), [3], [4], as complex matrices of the form

$$(1.1) \quad C := \begin{bmatrix} w & z \\ \bar{z} & \bar{w} \end{bmatrix},$$

where the bar $\bar{}$ indicates the complex conjugate of the quantity under the bar. The matrix C is very similar to a quaternion (invented by Hamilton, 1843), which has the complex representation (see v. d. Waerden, [16], p. 55)

$$Q := \begin{bmatrix} w & z \\ -\bar{z} & \bar{w} \end{bmatrix}.$$

In both cases, the products C_1C_2 , Q_1Q_2 of two coquaternions C_1, C_2 , and of two quaternions Q_1, Q_2 form a coquaternion, quaternion, respectively. Thus, both sets of matrices form a real algebra. This, in particular means that the *center* is in both cases the set of matrices $\alpha\mathbf{I}$, $\alpha \in \mathbb{R}$, where the center is the set of elements which commute with all elements and \mathbf{I} is the 2×2 identity matrix. There is one decisive difference between C and Q . The inverse of C is

$$C^{-1} := \frac{1}{|w|^2 - |z|^2} \begin{bmatrix} \bar{w} & -z \\ -\bar{z} & w \end{bmatrix}$$

and the inverse of Q is

$$Q^{-1} := \frac{1}{|w|^2 + |z|^2} \begin{bmatrix} \bar{w} & -z \\ \bar{z} & w \end{bmatrix}.$$

The inverse of C exists if and only if $|w|^2 - |z|^2 \neq 0$, whereas a quaternion Q has an inverse as long as $Q \neq 0$. Thus, the algebra of coquaternions has zero divisors, does not form a field, whereas the algebra of quaternions is free of zero divisors, it is a field, though not commutative.

* Institute of Chemical Technology, Prague, Department of Mathematics, Technická 5, 166 28 Prague 6, Czech Republic, janovskd@vscht.cz

† University of Hamburg, Faculty for Mathematics, Informatics, and Natural Sciences [MIN], Bundestraße 55, 20146 Hamburg, Germany, opfer@math.uni-hamburg.de

It is known, that a quaternion can also be represented by a real (4×4) matrix involving the four real numbers $\Re w, \Im w, \Re z, \Im z$, where \Re stands for real part and \Im stands for imaginary part of a complex number, see Gürlebeck and Sprössig, [7], p. 5.

THEOREM 1.1. *Put*

$$(1.2) \quad w = a_1 + a_2\mathbf{i}; \quad z = a_3 + a_4\mathbf{i},$$

and define the matrix

$$(1.3) \quad C_4 := \begin{bmatrix} a_1 & -a_2 & a_3 & a_4 \\ a_2 & a_1 & a_4 & -a_3 \\ a_3 & a_4 & a_1 & -a_2 \\ a_4 & -a_3 & a_2 & a_1 \end{bmatrix}.$$

Then, the set of all matrices of the type C_4 forms an algebra, and this algebra is isomorphic to the algebra of coquaternions.

Proof: Let C_4, \tilde{C}_4 be two matrices of type (1.3). Both have block structure,

$$C_4 = \begin{bmatrix} \mathbf{A}_1 & \mathbf{A}_2 \\ \mathbf{A}_2 & \mathbf{A}_1 \end{bmatrix}, \quad \mathbf{A}_1 := \begin{bmatrix} a_1 & -a_2 \\ a_2 & a_1 \end{bmatrix}, \quad \mathbf{A}_2 := \begin{bmatrix} a_3 & a_4 \\ a_4 & -a_3 \end{bmatrix},$$

$$\tilde{C}_4 = \begin{bmatrix} \mathbf{B}_1 & \mathbf{B}_2 \\ \mathbf{B}_2 & \mathbf{B}_1 \end{bmatrix}, \quad \mathbf{B}_1 := \begin{bmatrix} b_1 & -b_2 \\ b_2 & b_1 \end{bmatrix}, \quad \mathbf{B}_2 := \begin{bmatrix} b_3 & b_4 \\ b_4 & -b_3 \end{bmatrix},$$

hence,

$$C_4 \tilde{C}_4 = \begin{bmatrix} \mathbf{A}_1 \mathbf{B}_1 + \mathbf{A}_2 \mathbf{B}_2 & \mathbf{A}_1 \mathbf{B}_2 + \mathbf{A}_2 \mathbf{B}_1 \\ \mathbf{A}_1 \mathbf{B}_2 + \mathbf{A}_2 \mathbf{B}_1 & \mathbf{A}_1 \mathbf{B}_1 + \mathbf{A}_2 \mathbf{B}_2 \end{bmatrix},$$

$$\mathbf{A}_1 \mathbf{B}_1 + \mathbf{A}_2 \mathbf{B}_2 = \begin{bmatrix} a_1 b_1 - a_2 b_2 + a_3 b_3 + a_4 b_4 & -a_1 b_2 - a_2 b_1 + a_3 b_4 - a_4 b_3 \\ a_2 b_1 + a_1 b_2 + a_4 b_3 - a_3 b_4 & -a_2 b_2 + a_1 b_1 + a_4 b_4 + a_3 b_3 \end{bmatrix},$$

$$\mathbf{A}_1 \mathbf{B}_2 + \mathbf{A}_2 \mathbf{B}_1 = \begin{bmatrix} a_1 b_3 - a_2 b_4 + a_3 b_1 + a_4 b_2 & a_1 b_4 + a_2 b_3 - a_3 b_2 + a_4 b_1 \\ a_2 b_3 + a_1 b_4 + a_4 b_1 - a_3 b_2 & a_2 b_4 - a_1 b_3 - a_4 b_2 - a_3 b_1 \end{bmatrix}.$$

Thus, the product has the same structure as the matrix given in (1.3). If we compare the product of two coquaternions in the form given in (1.1) with the product $C_4 \tilde{C}_4$ where (1.2) is used, then, we see, that the products are the same. \square

Let $a_1, a_2, a_3, a_4 \in \mathbb{R}$ as in (1.2). The algebra of coquaternions is also isomorphic to the algebra of all real 2×2 matrices (see Lam, p. 52, [12]):

$$(1.4) \quad C_2 := a_1 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + a_2 \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} + a_3 \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} + a_4 \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

$$= \begin{bmatrix} a_1 + a_4 & a_2 + a_3 \\ -a_2 + a_3 & a_1 - a_4 \end{bmatrix} =: \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \end{bmatrix} \in \mathbb{R}^{2 \times 2}.$$

Given the four matrix elements $c_{11}, c_{12}, c_{21}, c_{22} \in \mathbb{R}$, the four components of a can be recovered by

$$a_1 = \frac{1}{2}(c_{11} + c_{22}), \quad a_2 = \frac{1}{2}(c_{12} - c_{21}), \quad a_3 = \frac{1}{2}(c_{12} + c_{21}), \quad a_4 = \frac{1}{2}(c_{11} - c_{22}).$$

The inverse of C_2 and of C_4 (given in (1.3)) can be computed easily by applying the formula (2.6) for a^{-1} . If we denote the four basis elements in the order of the equation (1.4) by $\mathbf{E}, \mathbf{I}, \mathbf{J}, \mathbf{K}$, then they obey the same multiplication rules as $1, \mathbf{i}, \mathbf{j}, \mathbf{k}$,

respectively, given in Table 2.1 of the next section. An algebra of this type, is also called a *split algebra*, in the current case the algebra of *split quaternions*, Lam, p. 58, [12].

That coquaternions nowadays are still useful, e. g. in physics is shown by Brody and Graefe, 2011, [2]. That paper also contains an overview over relevant properties of coquaternions and 42 references are quoted. There is another, very subtle investigation mainly on the analysis of coquaternions with application to physics by Frenkel and Libine, [6].

We will denote the set of real numbers, the set of complex numbers, the set of quaternions, the set of integers, the set of positive integers by $\mathbb{R}, \mathbb{C}, \mathbb{H}, \mathbb{Z}, \mathbb{N}$, respectively.

2. Coquaternions. In view of the preceding section, coquaternions may be regarded as elements of \mathbb{R}^4 of the form

$$a := a_1 + a_2\mathbf{i} + a_3\mathbf{j} + a_4\mathbf{k}, \quad a_1, a_2, a_3, a_4 \in \mathbb{R}$$

which we also abbreviate by $a = (a_1, a_2, a_3, a_4)$ and which obey the multiplication rules given in Table 2.1:

TABLE 2.1. Multiplication table for coquaternions, the red figures differ in sign from the corresponding table for quaternions.

	1	\mathbf{i}	\mathbf{j}	\mathbf{k}
1	1	\mathbf{i}	\mathbf{j}	\mathbf{k}
\mathbf{i}	\mathbf{i}	-1	\mathbf{k}	$-\mathbf{j}$
\mathbf{j}	\mathbf{j}	$-\mathbf{k}$	1	$-\mathbf{i}$
\mathbf{k}	\mathbf{k}	\mathbf{j}	\mathbf{i}	1

The algebra of coquaternions will be abbreviated by \mathbb{H}_{coq} . As elements of \mathbb{R}^4 we have $1 := (1, 0, 0, 0), \mathbf{i} := (0, 1, 0, 0), \mathbf{j} := (0, 0, 1, 0), \mathbf{k} := (0, 0, 0, 1)$, which we will abbreviate also by $\mathcal{N}_1, \mathcal{N}_2, \mathcal{N}_3, \mathcal{N}_4$, respectively. Let $a = (a_1, a_2, a_3, a_4), b = (b_1, b_2, b_3, b_4)$. For future purposes we note the following multiplication results:

$$(2.1) \quad \begin{aligned} ab = a\mathcal{N}_1b = & a_1b_1 - a_2b_2 + a_3b_3 + a_4b_4 \\ & + (a_1b_2 + a_2b_1 - a_3b_4 + a_4b_3)\mathbf{i} \\ & + (a_1b_3 - a_2b_4 + a_3b_1 + a_4b_2)\mathbf{j} \\ & + (a_1b_4 + a_2b_3 - a_3b_2 + a_4b_1)\mathbf{k}, \end{aligned}$$

$$(2.2) \quad \begin{aligned} a\mathbf{i}b = a\mathcal{N}_2b = & -a_2b_1 - a_1b_2 + a_4b_3 - a_3b_4 \\ & + (-a_2b_2 + a_1b_1 - a_4b_4 - a_3b_3)\mathbf{i} \\ & + (-a_2b_3 - a_1b_4 + a_4b_1 - a_3b_2)\mathbf{j} \\ & + (-a_2b_4 + a_1b_3 - a_4b_2 - a_3b_1)\mathbf{k}, \end{aligned}$$

$$(2.3) \quad \begin{aligned} a\mathbf{j}b = a\mathcal{N}_3b = & a_3b_1 - a_4b_2 + a_1b_3 + a_2b_4 \\ & + (a_3b_2 + a_4b_1 - a_1b_4 + a_2b_3)\mathbf{i} \\ & + (a_3b_3 - a_4b_4 + a_1b_1 + a_2b_2)\mathbf{j} \\ & + (a_3b_4 + a_4b_3 - a_1b_2 + a_2b_1)\mathbf{k}, \end{aligned}$$

$$(2.4) \quad \begin{aligned} a\mathbf{k}b = a\mathcal{N}_4b = & a_4b_1 + a_3b_2 - a_2b_3 + a_1b_4 \\ & + (a_4b_2 - a_3b_1 + a_2b_4 + a_1b_3)\mathbf{i} \\ & + (a_4b_3 + a_3b_4 - a_2b_1 + a_1b_2)\mathbf{j} \\ & + (a_4b_4 - a_3b_3 + a_2b_2 + a_1b_1)\mathbf{k}. \end{aligned}$$

A coquaternion of the form $a = (a_1, 0, 0, 0)$ will be called *real* and will also be abbreviated as a_1 . As we see from (2.1) (and former considerations), real coquaternions commute with all coquaternions. Let $a = (a_1, a_2, a_3, a_4)$ be a coquaternion. The first component, a_1 , of a , will be denoted by $\Re(a)$ and called *real part* of a . We define the *conjugate* of a in the notation \bar{a} or in the notation $\text{conj}(a)$ and abs_2 of a by

$$(2.5) \quad \bar{a} := (a_1, -a_2, -a_3, -a_4), \quad \text{abs}_2(a) := a_1^2 + a_2^2 - a_3^2 - a_4^2.$$

The quantity abs_2 may be negative, it is not the square of a norm. Let b be another coquaternion. There are the following rules:

$$a\bar{a} = \bar{a}a = \text{abs}_2(a), \quad \text{abs}_2(ab) = \text{abs}_2(ba) = \text{abs}_2(a)\text{abs}_2(b),$$

$$\overline{(ab)} = \bar{b}\bar{a}, \quad \Re(ab) = \Re(ba).$$

The coquaternion a will be called *singular* if $\text{abs}_2(a) = 0$. If a is nonsingular (= not singular = invertible), then

$$(2.6) \quad aa^{-1} = a^{-1}a = (1, 0, 0, 0) \text{ holds for } a^{-1} = \frac{\bar{a}}{\text{abs}_2(a)}.$$

Let the quaternion product (only in this section) be denoted by \star , then by comparing the coquaternion product (2.1) with the corresponding quaternion product \star , we see that

$$ab = a \star b + 2(a_3b_3 + a_4b_4) - 2(a_3b_4 - a_4b_3) \mathbf{i}.$$

The third and fourth component of ab and of $a \star b$ coincide. Thus, the coquaternions contain the complex numbers as subalgebra.

3. Linear mappings over \mathbb{R} in general. Let $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a linear mapping over \mathbb{R} . Then it is known that such a mapping can be represented by a real matrix of size $(m \times n)$. See Horn and Johnson, p. 5, [9]. In order to find this matrix, which we will denote, in this paper, by \mathbf{M} , we define a *column operator* col by

$$\text{col}(x) := \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix},$$

where x_1, x_2, \dots, x_n are the components of x . If it happens that x is a matrix, we put the columns of that matrix from the left to the right into one column in order to define col for that matrix. By evaluating L at $x \in \mathbb{R}^n$ and applying the col operator we obtain

$$(3.1) \quad \text{col}(L(x)) = \mathbf{M}\text{col}(x), \quad \mathbf{M} \in \mathbb{R}^{m \times n}, \text{col}(x) \in \mathbb{R}^n,$$

where \mathbf{M} is unknown so far. Let e_j be the standard unit vectors in \mathbb{R}^n , $j = 1, 2, \dots, n$. If we put $x := e_j$ we obtain

$$(3.2) \quad \text{col}(L(e_j)) = \mathbf{M}\text{col}(e_j) = \begin{bmatrix} \mu_{1j} \\ \mu_{2j} \\ \vdots \\ \mu_{mj} \end{bmatrix}, \quad j = 1, \dots, n,$$

where μ_{ij} are the elements of the matrix \mathbf{M} and the right hand side of (3.2) represents the j th column of \mathbf{M} . Hence, the matrix \mathbf{M} is completely known by the n values $L(e_j)$. We note, that \mathbf{M} will be integer if the values of L are integer.

A typical, nontrivial example is the mapping $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$ defined by

$$(3.3) \quad L(\mathbf{X}) = \mathbf{A}\mathbf{X}\mathbf{B}, \quad \mathbf{A} \in \mathbb{R}^{p \times q}, \quad \mathbf{B} \in \mathbb{R}^{r \times s}, \quad \mathbf{X} \in \mathbb{R}^{q \times r},$$

where in this case we have $m = ps$, $n = qr$. In order to find the corresponding matrix we have to set the j th element of \mathbf{X} (counted columnwise from the left) to be one and all other elements of \mathbf{X} to be zero, $j = 1, 2, \dots, qr$, and compute the corresponding $\text{col}(L(\mathbf{X}))$. This gives the j th column of the wanted matrix \mathbf{M} . What comes out is the well known Kronecker (or tensor) product $\mathbf{M} = \mathbf{B}^T \otimes \mathbf{A}$, a matrix of size $(ps \times qr)$. The notation \mathbf{B}^T stands for the *transposed matrix* of \mathbf{B} . For details see Horn and Johnson, pp. 242, 254, [8].

4. Linear systems in coquaternions. Linear equations in coquaternions are formally similar to linear equations in quaternions. An investigation of linear equations in quaternions exists by the current authors, [10], [11] and by Niven, [13]. A linear system will always be a linear system over \mathbb{R} . Linearity with respect to \mathbb{C} or to \mathbb{H}_{coq} is in general not granted. A linear system in n coquaternions x_k , $k = 1, 2, \dots, n$, and m equations will be defined as follows: Let

$$(4.1) \quad l_j(u) := \sum_{k=1}^{K_j} a_k^{(j)} u b_k^{(j)}, \quad j = 1, 2, \dots, mn$$

be an arbitrary set of mn linear, coquaternionic mappings in one coquaternionic variable u , where $a_k^{(j)}, b_k^{(j)}$, $k = 1, 2, \dots, K_j$ are given coquaternions, and K_j are given, positive integers, $j = 1, 2, \dots, mn$. A system in n coquaternionic variables x_k , $k = 1, 2, \dots, n$ and m equations will then be defined by

$$\begin{aligned} L_1(x) &:= \sum_{j=1}^n l_j(x_j), \\ L_2(x) &:= \sum_{j=1}^n l_{j+n}(x_j), \\ &\vdots \\ L_m(x) &:= \sum_{j=1}^n l_{j+(m-1)n}(x_j), \end{aligned}$$

where $x \in \mathbb{R}^{4n}$ consists of one column composed out of x_1, x_2, \dots, x_n . The mapping $L : \mathbb{R}^{4n} \rightarrow \mathbb{R}^{4m}$ will finally be defined by

$$(4.2) \quad L(x) := \begin{bmatrix} L_1(x) \\ L_2(x) \\ \vdots \\ L_m(x) \end{bmatrix}.$$

L is not a linear mapping over \mathbb{C} or \mathbb{H}_{coq} . However, L is a linear mapping over \mathbb{R} because real coquaternions commute with arbitrary coquaternions. Hence, a matrix \mathbf{M} as in (3.1), (3.2) exists.

In order to find the matrix \mathbf{M} which represents L , we put $x = e_j$, where the e_j represent the standard unit vectors in \mathbb{R}^{4n} , $j = 1, 2, \dots, 4n$. Then $\text{col}(L(e_j))$ is the j th column of the wanted matrix. The next lemma will be useful.

LEMMA 4.1. *Let $1 \leq j \leq 4n$ be given. Then, the following equation including its restrictions*

$$(4.3) \quad 4(\kappa - 1) + r = j, \quad 1 \leq \kappa \leq n, \quad 1 \leq r \leq 4$$

determines κ and r uniquely by

$$(4.4) \quad 0 \leq 4\kappa - j \leq 3, \quad r = j - 4(\kappa - 1).$$

Proof: If we solve (4.3) for r we obtain $r = j - 4(\kappa - 1) = j - 4\kappa + 4$ and the given restriction for r implies the first part of (4.4) and the first part admits exactly one solution κ with the restriction given in (4.3). For this reason, the second equation in (4.4) (which coincides with the first equation in (4.3)) also has a unique solution. \square

COROLLARY 4.2. *Let L be given as in (4.2). Then the j th column ($1 \leq j \leq 4n$) of the matrix \mathbf{M} representing L is*

$$(4.5) \quad L(e_j) := \begin{bmatrix} L_1(e_j) = l_\kappa(\mathcal{N}_r) \\ L_2(e_j) = l_{\kappa+n}(\mathcal{N}_r) \\ \vdots \\ L_m(e_j) = l_{\kappa+(m-1)n}(\mathcal{N}_r) \end{bmatrix}$$

where κ and r are determined by j applying Lemma 4.1. The first component of the unit vector e_j is the r th coordinate, $r \in \{1, 2, 3, 4\}$, of the κ th coquaternion, $\kappa \in \{1, 2, \dots, n\}$.

The task of solving a linear, coquaternionic system $L(x) = \gamma$, where γ consists of m given coquaternions, can be solved as follows. Compute the $(4m \times 4n)$ matrix \mathbf{M} which represents L and solve the real matrix equation $\mathbf{M}\text{col}(x) = \text{col}(\gamma)$ by standard techniques.

EXAMPLE 4.3. Let $m = n = 3$ and

$$\begin{aligned} l_1(x) &:= a_1^{(1)}xb_1^{(1)} + a_2^{(1)}xb_2^{(1)} + a_3^{(1)}xb_3^{(1)}, \quad l_2(x) := a_1^{(2)}xb_1^{(2)} + a_2^{(2)}xb_2^{(2)}, \quad l_3(x) := a_1^{(3)}xb_1^{(3)}, \\ l_4(x) &:= a_1^{(4)}xb_1^{(4)}, \quad l_5(x) := a_1^{(5)}xb_1^{(5)} + a_2^{(5)}xb_2^{(5)}, \quad l_6(x) := a_1^{(6)}xb_1^{(6)} + a_2^{(6)}xb_2^{(6)} + a_3^{(6)}xb_3^{(6)}, \\ l_7(x) &:= a_1^{(7)}xb_1^{(7)} + a_2^{(7)}xb_2^{(7)}, \quad l_8(x) := a_1^{(8)}xb_1^{(8)} + a_2^{(8)}xb_2^{(8)}, \quad l_9(x) := a_1^{(9)}xb_1^{(9)} + a_2^{(9)}xb_2^{(9)}, \end{aligned}$$

where the needed coefficients $a_k^{(j)}, b_k^{(j)}$ are defined in Table 4.4.

TABLE 4.4. Table of coquaternionic coefficients for l_1 to l_9

j	$a_1^{(j)}$	$b_1^{(j)}$	$a_2^{(j)}$	$b_2^{(j)}$	$a_3^{(j)}$	$b_3^{(j)}$
1	(0, 0, -1, 1)	(-1, 1, 2, 0)	(0, 1, -1, -1)	(-1, 2, 2, -1)	(0, -1, 1, 1)	(-1, 2, 2, -1)
2	(2, 0, 1, 0)	(-1, 2, 1, 1)	(-2, 0, 1, 0)	(-1, 0, 1, 2)		
3	(1, -1, 1, -2)	(0, 1, 2, 1)				
4	(0, 1, 0, 2)	(2, 1, 0, 0)				
5	(-1, 1, 2, 2)	(0, -1, 2, 1)	(0, 0, -1, 2)	(2, -1, 1, 1)		
6	(-2, 0, 1, -1)	(0, -1, 0, 1)	(-1, 2, 1, 0)	(2, 1, -2, 1)	(0, 1, 1, 0)	(0, 0, -2, 1)
7	(-1, 1, 1, 0)	(0, 1, 2, 1)	(0, 0, 0, 0)	(0, 1, -1, -2)		
8	(-2, 1, 1, 1)	(1, 1, 2, 0)	(-2, 0, 2, 1)	(2, 0, 0, -1)		
9	(0, 0, -2, 2)	(0, 1, 2, 2)	(-2, 0, 1, 0)	(-1, 1, -2, 0)		

The entries of the Table 4.4 are randomly generated integers in $[-2, 2]$. This is the reason for the occurrence of a zero element. The matrix \mathbf{M} which in this case corresponds to L , defined in (4.5), is

$$\mathbf{M} = \begin{bmatrix} 3 & -5 & -1 & 2 & 2 & -7 & -2 & 0 & 1 & -6 & 3 & 4 \\ -9 & -3 & 6 & -5 & 1 & -2 & 4 & 2 & -4 & 1 & -2 & -1 \\ 3 & 0 & -3 & 1 & -2 & 0 & 2 & 7 & 1 & 0 & 3 & -2 \\ 4 & 3 & -3 & 3 & -4 & 2 & -1 & -2 & -2 & 5 & -4 & -5 \\ -1 & -2 & -2 & 4 & 8 & 4 & 1 & 0 & -7 & -6 & 8 & 5 \\ 2 & -1 & 4 & 2 & 6 & -6 & 6 & -3 & 4 & -1 & -1 & 2 \\ 2 & 4 & 1 & -2 & -9 & 4 & -2 & 8 & 0 & 7 & -5 & -2 \\ 4 & -2 & 2 & 1 & 6 & 3 & 6 & 0 & -11 & -2 & 4 & 5 \\ -1 & 2 & -3 & 0 & -6 & -1 & -8 & 0 & 4 & 6 & -7 & -3 \\ 0 & 3 & -2 & -3 & -7 & -4 & -8 & 0 & 1 & 0 & -3 & -1 \\ -1 & 4 & -3 & -2 & -10 & -2 & -10 & -1 & -3 & 3 & -4 & 2 \\ -2 & -1 & 0 & 1 & -2 & 2 & 1 & -4 & 3 & -5 & -2 & 8 \end{bmatrix}.$$

Let $x_1 = (0, 0, 1, 0)$, $x_2 = (-1, 0, 0, 2)$, $x_3 = (0, 1, 1, -1)$ and $x = \text{col}(x_1, x_2, x_3)$. Then,

$$\text{col}(L(x)) := \mathbf{M}x = (-10, 9, 18, 3; -13, -12, 30, -7; 5, 3, 2, -21)^T.$$

4.1. Linear systems in coquaternionic matrices. The general case of a linear system in n unknowns and m equations does also contain the matrix case

$$(4.6) \quad \mathbf{A}\mathbf{x} = \mathbf{b}, \quad \mathbf{A} \in \mathbb{H}_{\text{coq}}^{m \times n}, \quad \mathbf{b} \in \mathbb{H}_{\text{coq}}^{m \times 1}, \quad \mathbf{x} \in \mathbb{H}_{\text{coq}}^{n \times 1},$$

but not vice versa.

EXAMPLE 4.5. Let

$$\mathbf{A} := \begin{bmatrix} 1 & \mathbf{i} \\ \mathbf{j} & \mathbf{k} \end{bmatrix} \Rightarrow \mathbf{A}^{-1} = \frac{1}{2} \begin{bmatrix} 1 & \mathbf{j} \\ -\mathbf{i} & \mathbf{k} \end{bmatrix}, \quad \mathbf{A}^T = \begin{bmatrix} 1 & \mathbf{j} \\ \mathbf{i} & \mathbf{k} \end{bmatrix}.$$

If we multiply the second row of \mathbf{A} from the right by \mathbf{j} we obtain the first row. Thus, the right row rank of \mathbf{A} is one, and if we multiply the first column from the left by \mathbf{i} we obtain the second column. Thus, the left column rank of \mathbf{A} is also one. Let us compute the right column rank of \mathbf{A} . To this purpose let

$$\begin{pmatrix} 1 \\ \mathbf{j} \end{pmatrix} \alpha + \begin{pmatrix} \mathbf{i} \\ \mathbf{k} \end{pmatrix} \beta = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \quad \alpha, \beta \in \mathbb{H}_{\text{coq}}.$$

If we multiply the first equation from the left by \mathbf{j} we obtain $\mathbf{j}\alpha - \mathbf{k}\beta = 0$. If we add this equation to the second equation we obtain $2\mathbf{j}\alpha = 0$, and if we subtract this equation from the second equation we obtain $2\mathbf{k}\beta = 0$. Thus, $\alpha = \beta = 0$ and the right column rank is two. In a similar way one can show that the left row rank is also two. There is the following well known Theorem: Let $\mathbf{M} \in \mathbb{H}_{\text{coq}}^{m \times n}$. Then, the left row rank coincides with the right column rank and the right row rank is equal to the left column rank. If $m = n$ and the right column rank of \mathbf{M} is n , then \mathbf{M} is nonsingular, which means that there is a matrix \mathbf{M}^{-1} such that of $\mathbf{M}^{-1}\mathbf{M} = \mathbf{M}\mathbf{M}^{-1} = \mathbf{I}$, where \mathbf{I} is the $(n \times n)$ identity matrix. Thus, \mathbf{A} is nonsingular, whereas \mathbf{A}^T is singular. This example was also used by Zhang, [17], for quaternions.

Rather than specializing the general case to the matrix case, we directly treat the matrix case (4.6) with the means we have already derived. If we apply the col operator to (4.6) we obtain

$$(4.7) \quad \text{col}(\mathbf{A}\mathbf{x}) = \mathbf{M}\text{col}(\mathbf{x}) = \text{col}(\mathbf{b}), \quad \mathbf{M} \in \mathbb{R}^{4m \times 4n}$$

and determine \mathbf{M} by replacing $\text{col}(\mathbf{x})$ with e_j , the j th standard unit vector in \mathbb{R}^{4n} , $j = 1, 2, \dots, 4n$. Let $j = 1 \pmod 4$, $j \leq 4n - 3$, $k = 1, 2, 3, 4$. Then, the $(j+k-1)$ th column of \mathbf{M} is

$$(4.8) \quad \mathbf{M}_{(j+k-1)} = \begin{bmatrix} \text{col}(a_{1,(j-1)/4+1}\mathcal{N}_k) \\ \text{col}(a_{2,(j-1)/4+1}\mathcal{N}_k) \\ \vdots \\ \text{col}(a_{m,(j-1)/4+1}\mathcal{N}_k) \end{bmatrix}.$$

Each entry, $\text{col}(a_{\ell,(j-1)/4+1}\mathcal{N}_k)$, is a real 4×1 vector, $\ell = 1, 2, \dots, m$, where we denote the j th column of \mathbf{M} by $\mathbf{M}_{(j)}$. Thus, \mathbf{M} of (4.7) is given in (4.8). For \mathbf{A} from Example 4.5 we have

$$\mathbf{M} = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & -1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & -1 & 0 & 0 & 1 & 0 & 0 & 0 \end{bmatrix}.$$

There are many further questions related to coquaternionic matrices, for instance, what kind of decompositions exist. This is a topic for another investigation.

4.2. The Kronecker product for coquaternions. The *coquaternionic Kronecker product* is the (real) matrix \mathbf{M} which represents the linear mapping (over \mathbb{R}) $L: \mathbb{H}_{\text{coq}}^{q \times r} \rightarrow \mathbb{H}_{\text{coq}}^{p \times s}$ defined by

$$(4.9) \quad L(\mathbf{X}) = \mathbf{A}\mathbf{X}\mathbf{B}, \quad \mathbf{A} \in \mathbb{H}_{\text{coq}}^{p \times q}, \quad \mathbf{B} \in \mathbb{H}_{\text{coq}}^{r \times s}, \quad \mathbf{X} \in \mathbb{H}_{\text{coq}}^{q \times r}.$$

It is denoted by $\mathbf{M}(\mathbf{A}, \mathbf{B})$, for short only \mathbf{M} . Since each element in \mathbf{X} has 4 real entries, $\text{col}(\mathbf{X}) \in \mathbb{R}^{4qr}$. Furthermore, we have $\text{col}(L(\mathbf{X})) \in \mathbb{R}^{4ps}$ which implies $\mathbf{M} \in \mathbb{R}^{4ps \times 4qr}$. The coquaternionic Kronecker product differs from the standard Kronecker or tensor product which is $\mathbf{B}^T \otimes \mathbf{A}$. See also (3.3), on p. 5 and the corresponding remarks. In order to find \mathbf{M} , we enumerate the elements of \mathbf{X} by $\ell = 1, 2, \dots, qr$, using the columns as ordering system such that (to mention an example) the first element of the second column will have the number $\ell = q + 1$. We denote the so numbered elements of \mathbf{X} by $\mathbf{X}(1), \mathbf{X}(2), \dots, \mathbf{X}(qr)$. Such an enumeration technique is used in some programming languages in addition to the conventional enumeration $\mathbf{X}_{J,K}$, which denotes the K th element in row J . We define the matrix $\mathbf{X}^{(\ell,k)}$ by putting

$$(4.10) \quad \mathbf{X}(\rho) = \begin{cases} \mathcal{N}_k & \text{for } \rho = \ell, \\ 0 & \text{for } \rho \neq \ell, \end{cases} \quad \ell = 1, 2, \dots, qr, \quad k = 1, 2, 3, 4.$$

In other words, $\mathbf{X}^{(\ell,k)}$ is the matrix which contains \mathcal{N}_k at position ℓ , and contains otherwise zero elements. The corresponding columns of \mathbf{M} are then

$$(4.11) \quad \mathbf{M}_{(4(\ell-1)+k)} := \text{col}(\mathbf{A}\mathbf{X}^{(\ell,k)}\mathbf{B}), \quad \ell = 1, 2, \dots, qr, \quad k = 1, 2, 3, 4.$$

We will describe an alternative technique to find \mathbf{M} . Let us assume that \mathbf{X} contains only zeros, with the exception of the element $\mathbf{X}_{J,K}$ which contains one of the four unit elements \mathcal{N}_k . Then,

$$(4.12) \quad \mathbf{M}_{4((K-1)q+(J-1))+k} = \text{col}(\mathbf{A}\mathbf{X}\mathbf{B}) = \text{col} \left(\begin{array}{c} \begin{array}{cccccc} 1 & \dots & K-1 & K & K+1 & r \\ 0 & \dots & 0 & a_{1J} & 0 & \dots & 0 \\ 0 & \dots & 0 & a_{2J} & 0 & \dots & 0 \\ 0 & \dots & 0 & \vdots & 0 & \dots & 0 \\ 0 & \dots & 0 & a_{pJ} & 0 & \dots & 0 \end{array} \\ \begin{array}{cccccc} 0 & \dots & 0 & \dots & 0 \\ \mathcal{N}_k b_{K1} & \mathcal{N}_k b_{K2} & \dots & \dots & \mathcal{N}_k b_{Ks} \\ 0 & \dots & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & \dots & 0 & \dots & 0 \end{array} \end{array} \right).$$

THEOREM 4.6. *Let $K \in [1, r], J \in [1, q], k \in [1, 4]$ be fixed. See (4.9). The coquaternionic Kronecker product $\mathbf{M}(\mathbf{A}, \mathbf{B})$ in column $[4((K-1)q + (J-1)) + k]$ is given by the dyadic coquaternionic product $\text{col}(\mathbf{A}(:, J)(\mathcal{N}_k \mathbf{B}(K, :)))$.*

Proof: Follows from the representation (4.12). \square

EXAMPLE 4.7. We will treat an example of the form

$$\mathbf{C} = \mathbf{A}\mathbf{X}\mathbf{B} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \begin{bmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \end{bmatrix} \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix}$$

where we use the data from Example 5.3 in [10]:

$$a_{11} = (0, 2, 2, 0), \quad a_{12} = (4, 5, -1, -5), \quad a_{21} = (0, 2, 2, -1), \quad a_{22} = (-3, 3, -3, 2),$$

$$b_{11} = (0, 4, -5, -4), \quad b_{12} = (-2, 2, 1, -4), \quad b_{21} = (-3, -5, 2, -1), \quad b_{22} = (4, 3, -2, 3),$$

$$x_{11} = (1, 1, 1, 1), \quad x_{12} = (1, 2, 1, 2), \quad x_{21} = (2, 1, 2, 1), \quad x_{22} = (2, 2, 2, 2).$$

These data determine $\mathbf{C} = \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \end{bmatrix}$ as

$$c_{11} = (28, -1, 44, -65), \quad c_{12} = (-58, -13, 2, 35),$$

$$c_{21} = (44, -92, 132, -12), \quad c_{22} = (-76, -79, 32, 89).$$

An application of (4.11) or of Theorem 4.6 yields the Kronecker product

$$\mathbf{M}(\mathbf{A}, \mathbf{B}) = \begin{bmatrix} -18 & 8 & -8 & 18 & 5 & 5 & -20 & 5 \\ 8 & 2 & -2 & -8 & 37 & -45 & -13 & -60 \\ 8 & 2 & -2 & -8 & -20 & 45 & 5 & 45 \\ -18 & 8 & -8 & 18 & -37 & 20 & 13 & 35 \\ -14 & 13 & -4 & 18 & -5 & -10 & -5 & 15 \\ 13 & -2 & -2 & -12 & -34 & -19 & -39 & 11 \\ 4 & 2 & -6 & -3 & 35 & 15 & 35 & -10 \\ -18 & 12 & -3 & 22 & 9 & 19 & 14 & -11 \\ -2 & 12 & -12 & 2 & 1 & -7 & -4 & -13 \\ 4 & -6 & 6 & -4 & -11 & -37 & 29 & -28 \\ 4 & -6 & 6 & -4 & 16 & 23 & -19 & 17 \\ -2 & 12 & -12 & 2 & 1 & 32 & -19 & 23 \\ 2 & 11 & -10 & 4 & -11 & 2 & -13 & -1 \\ 3 & -10 & 8 & -6 & -22 & 11 & -19 & -17 \\ 2 & -4 & 2 & -5 & 19 & -13 & 17 & 14 \\ 0 & 14 & -13 & 6 & 17 & -1 & 14 & 7 \end{bmatrix} \dots$$

$$\begin{bmatrix} 14 & 8 & -8 & -14 & 16 & 24 & -19 & 6 \\ -4 & 6 & -6 & 4 & -46 & 10 & 34 & 25 \\ -4 & 6 & -6 & 4 & 41 & 4 & -44 & -14 \\ 14 & 8 & -8 & -14 & 16 & -15 & -4 & -30 \\ 15 & 6 & -13 & -11 & 16 & -5 & 10 & 6 \\ -6 & 5 & -3 & 9 & 7 & 32 & 12 & -28 \\ 1 & 9 & -7 & 2 & -4 & -30 & -10 & 31 \\ 17 & 3 & -10 & -13 & -12 & -2 & -17 & -2 \\ -10 & -14 & 14 & 10 & -12 & -19 & 18 & -1 \\ 2 & -2 & 2 & -2 & 45 & 14 & -45 & -4 \\ 2 & -2 & 2 & -2 & -42 & -19 & 48 & -1 \\ -10 & -14 & 14 & 10 & -15 & -4 & 15 & 14 \\ -13 & -12 & 17 & 6 & -9 & 2 & -3 & -4 \\ 4 & 1 & -2 & -5 & 8 & -33 & 2 & 33 \\ -1 & -6 & 5 & 0 & -9 & 32 & -3 & -34 \\ -14 & -11 & 16 & 7 & 2 & 3 & 8 & -3 \end{bmatrix} \in \mathbb{R}^{16 \times 16}.$$

A further test shows that $\mathbf{M}\text{col}(\mathbf{X}) = \text{col}(\mathbf{C})$ for \mathbf{X} chosen in the beginning of this example.

Another interesting example is Sylvester's equation [15, Sylvester, 1884]:

$$(4.13) \quad \mathbf{A}\mathbf{X} + \mathbf{X}\mathbf{B} = \mathbf{C}, \quad \mathbf{A} \in \mathbb{H}_{\text{coq}}^{m \times m}, \mathbf{B} \in \mathbb{H}_{\text{coq}}^{n \times n}, \mathbf{C}, \mathbf{X} \in \mathbb{H}_{\text{coq}}^{m \times n}$$

$$\Rightarrow \text{col}(\mathbf{A}\mathbf{X} + \mathbf{X}\mathbf{B}) = (\mathbf{M}(\mathbf{A}, \mathbf{I}) + \mathbf{M}(\mathbf{I}, \mathbf{B}))\text{col}(\mathbf{X}) = \text{col}(\mathbf{C}).$$

It may be that the linear function L has the form

$$L(\mathbf{X}) = \mathbf{A}\mathbf{X}^T\mathbf{B}, \quad \mathbf{A} \in \mathbb{H}_{\text{coq}}^{p \times q}, \mathbf{B} \in \mathbb{H}_{\text{coq}}^{r \times s}, \mathbf{X} \in \mathbb{H}_{\text{coq}}^{r \times q}.$$

In order to find the corresponding Kronecker product for this case one has to change the order in which the matrix elements are counted. The matrix elements of \mathbf{X} in formula (4.10) have to be counted row-wise.

Let \mathbf{A}, \mathbf{B} be two coquaternionic matrices. It should be noted, that the relation $(\mathbf{A}\mathbf{B})^T = \mathbf{B}^T\mathbf{A}^T$ is in general not true. Let us denote by $\text{conj}(\mathbf{A})$ the matrix where all elements of \mathbf{A} have been changed to the corresponding conjugate element. Then, the relation $\text{conj}(\mathbf{A}\mathbf{B}) = \text{conj}(\mathbf{A})\text{col}(\mathbf{B})$ is in general also not true. However, $(\mathbf{A}\mathbf{B})^* = \mathbf{B}^*\mathbf{A}^*$, is true, where $\mathbf{A}^* := \text{conj}(\mathbf{A}^T)$.

It should be pointed out that the investigations of this paper are not restricted to equations in coquaternions. With the same technique one could solve linear systems of equations in *tessarines* or in *cotessarines*, algebras defined in \mathbb{R}^4 and also introduced by Cockle, [3]. Even more general algebras (finite dimensional, real, associative algebras) would allow the application of the same technique. See Abian, Drozd and Kirichenko, Pierce [1, 5, 14].

5. The one dimensional case. For $n = m = 1$ the system (4.2) and (4.1) specialize to one equation

$$(5.1) \quad L(x) := \sum_{k=1}^K a_k x b_k, \quad \text{where } a_k, b_k, x \text{ are coquaternions.}$$

The simplest cases are

$$(i) L(x) = ax, \quad (ii) L(x) = xb, \quad (iii) L(x) = axb,$$

We denote the real 4×4 matrices for the three cases, respectively, by

$${}_a\mathbf{M}, \quad \mathbf{M}_b, \quad {}_a\mathbf{M}_b.$$

Let $a = (a_1, a_2, a_3, a_4), b = (b_1, b_2, b_3, b_4)$. Then the three matrices are

$$(5.2) \quad {}_a\mathbf{M} := (a\mathcal{N}_1, a\mathcal{N}_2, a\mathcal{N}_3, a\mathcal{N}_4) = \begin{bmatrix} a_1 & -a_2 & a_3 & a_4 \\ a_2 & a_1 & a_4 & -a_3 \\ a_3 & a_4 & a_1 & -a_2 \\ a_4 & -a_3 & a_2 & a_1 \end{bmatrix},$$

$$(5.3) \quad \mathbf{M}_b := (\mathcal{N}_1b, \mathcal{N}_2b, \mathcal{N}_3b, \mathcal{N}_4b) = \begin{bmatrix} b_1 & -b_2 & b_3 & b_4 \\ b_2 & b_1 & -b_4 & b_3 \\ b_3 & -b_4 & b_1 & b_2 \\ b_4 & b_3 & -b_2 & b_1 \end{bmatrix},$$

$$(5.4) \quad {}_a\mathbf{M}_b := [ab, a\mathcal{N}_2b, a\mathcal{N}_3b, a\mathcal{N}_4b],$$

where the four columns are already defined in (2.1) to (2.4). Matrix ${}_a\mathbf{M}$ is identical with the matrix C_4 , defined in (1.3) and this is clear because C_4 represents the coquaternion $a = (a_1, a_2, a_3, a_4)$ and the mapping L is defined by $L(x) = ax$. A real, square matrix is named *singular* if it is not invertible, or in other words if its determinant vanishes.

THEOREM 5.1. *Matrix ${}_a\mathbf{M}$ is singular if and only if a is singular. Matrix \mathbf{M}_b is singular if and only if b is singular. Matrix ${}_a\mathbf{M}_b$ is singular if a or b is singular.*

Proof: Let $L(x) = ax$ and $a \neq 0$. The equation $ax = c$ has clearly a unique solution if a is nonsingular. If a is singular, we multiply by \bar{a} and obtain $0 = \bar{a}ax = \bar{a}c$, which does not have a unique solution. The other cases are similar. \square

EXAMPLE 5.2. We apply the technique described in Section 3 to

$$\begin{aligned} L(x) &= a_1xb_1 + a_2xb_2 + a_3xb_3, \quad \text{where} \\ a_1 &= (1, 1, 0, 1), \quad b_1 = (1, -1, -1, 0), \\ a_2 &= (1, -1, -1, 2), \quad b_2 = (2, -1, 1, 1), \\ a_3 &= (-1, -2, 0, 2), \quad b_3 = (0, -1, 0, 0). \end{aligned}$$

From these data we obtain

$$\begin{aligned} a_1b_1 &= (2, -1, -2, 0), \quad a_1\mathbf{i}b_1 = (-1, 2, 2, 0), \\ a_1\mathbf{j}b_1 &= (0, 0, 0, 1), \quad a_1\mathbf{k}b_1 = (2, -2, -3, 0), \\ a_2b_2 &= (2, 0, -2, 3), \quad a_2\mathbf{i}b_2 = (6, 0, 3, 6), \\ a_2\mathbf{j}b_2 &= (0, 3, 0, 0), \quad a_2\mathbf{k}b_2 = (7, 0, 2, 6), \\ a_3b_3 &= (-2, 1, -2, 0), \quad a_3\mathbf{i}b_3 = (-1, -2, 0, 2), \\ a_3\mathbf{j}b_3 &= (2, 0, 2, -1), \quad a_3\mathbf{k}b_3 = (0, -2, 1, 2), \end{aligned}$$

and the matrix \mathbf{M} which represents the linear mapping L is

$$\mathbf{M} = \begin{bmatrix} 2 & 4 & 2 & 9 \\ 0 & 0 & 3 & -4 \\ -6 & 5 & 2 & 0 \\ 3 & 8 & 0 & 8 \end{bmatrix}.$$

In order to find the four columns of \mathbf{M} , one has to compute

$$\text{col} \left(\sum_{j=1}^3 a_j\mathcal{N}_k b_j \right), \quad k = 1, 2, 3, 4.$$

5.1. Sylvester's equation in coquaternions. Sylvester's equation (see [15]) in coquaternions, a special case of (4.13) and of (5.1), reads

$$(5.5) \quad L(x) := ax + xb = c, \quad a, b, c, x \in \mathbb{H}_{\text{coq}},$$

where we will put $a := (a_1, a_2, a_3, a_4)$, $b := (b_1, b_2, b_3, b_4) \in \mathbb{H}_{\text{coq}}$. It is clear that the matrix corresponding to this system is

$$(5.6) \quad \mathbf{M} := {}_a\mathbf{M} + \mathbf{M}_b = \begin{bmatrix} s_1 & -s_2 & s_3 & s_4 \\ s_2 & s_1 & d_4 & -d_3 \\ s_3 & d_4 & s_1 & -d_2 \\ s_4 & -d_3 & d_2 & s_1 \end{bmatrix},$$

where ${}_a\mathbf{M}, \mathbf{M}_b$ are defined in (5.2), (5.3) and where

$$s_j := a_j + b_j, \quad d_j := a_j - b_j, \quad j = 1, 2, 3, 4.$$

Sylvester's equation will be called *singular* if the matrix \mathbf{M} defined in (5.6) is singular. We observe that $d_1 := a_1 - b_1$ is not occurring in (5.6).

EXAMPLE 5.3. Let $a = (1, 1, 1, 2)$, $b = (1, -2, 2, 1)$. Then,

$$\mathbf{M} := \begin{bmatrix} 2 & 1 & 3 & 3 \\ -1 & 2 & 1 & 1 \\ 3 & 1 & 2 & -3 \\ 3 & 1 & 3 & 2 \end{bmatrix}.$$

In this case a is nonsingular, b is singular, and \mathbf{M} is, nevertheless, nonsingular.

LEMMA 5.4. *The determinant of \mathbf{M} , defined in (5.6) is*

$$(5.7) \quad \det(\mathbf{M}) = s_1^2 (s_1^2 + s_2^2 + d_2^2 - (s_3^2 + d_3^2) - (s_4^2 + d_4^2)) + (-s_2d_2 + s_3d_3 + s_4d_4)^2.$$

Proof: Apply the expansion formula to (5.6) and collect terms. \square

The singularity of Sylvester's equation, expressed by the singularity of the matrix \mathbf{M} , is characterized in the next theorem.

THEOREM 5.5. *The matrix \mathbf{M} , defined in (5.6) is singular if and only if either*

$$s_1 := a_1 + b_1 = 0 \text{ and } -\text{abs}_2(a) + \text{abs}_2(b) = 0, \quad \text{or } s_1 \neq 0 \text{ and}$$

$$s_1^2 (-d_1^2 + 2(\text{abs}_2(a) + \text{abs}_2(b))) + (-\text{abs}_2(a) + \text{abs}_2(b) + s_1d_1)^2 = 0,$$

where the definition of abs_2 is given in (2.5).

Proof: Use $s_j^2 + d_j^2 = 2(a_j^2 + b_j^2)$ and $s_jd_j = a_j^2 - b_j^2$, $j = 1, 2, 3, 4$ and insert this into (5.7) of Lemma 5.4. \square

An example where the second part of Theorem 5.5 applies is $a = (2, 1, 1, 2)$, $b = (1, 2, 2, 5)$. In this case a is singular, b is nonsingular, but \mathbf{M} is singular.

COROLLARY 5.6. *Let both, $a, b \in \mathbb{H}_{\text{coq}}$ be singular. Then, \mathbf{M} defined in (5.6) is singular.*

Proof: The assumptions are $\text{abs}_2(a) = \text{abs}_2(b) = 0$. The singularity of \mathbf{M} follows from one of the conditions of Theorem 5.5. \square

Conversely, one cannot say, that nonsingular a, b imply a nonsingular matrix \mathbf{M} . A necessary, but not sufficient condition for \mathbf{M} being nonsingular is, that at least a or b is nonsingular.

EXAMPLE 5.7. Let $a = (a_1, a_2, a_3, a_4) = (0, 0, 1, 0) = \mathbf{j}$, $b = (b_1, b_2, b_3, b_4) = (0, 0, 0, 1) = \mathbf{k}$. Then, $s_1 = a_1 + b_1 = 0$, $\text{abs}_2(a) = -1$, $\text{abs}_2(b) = -1$, thus, both a, b are nonsingular, but the first condition of Theorem 5.5 is satisfied, and, hence \mathbf{M} , defined in (5.6), and Sylvester's equation (5.5) are singular.

Acknowledgments. The research of the second mentioned author was supported by the German Science Foundation, DFG, GZ: OP 33/19-1. We would like to thank the referee for presenting a very detailed report, which resulted in many improvements of our paper. There is no warranty, that web pages which occur in the list of references will stay unchanged or remain existing.

REFERENCES

- [1] A. ABIAN, *Linear associative algebras*, Pergamon Press, New York, 1971, 166 pp.
- [2] D. C. BRODY, E.-M. GRAEFE, *On complexified mechanics and coquaternions*, J. Phys. A 44 (2011) 072001, 9 pp. Also: arXiv:1012.0757v2 [math-ph]
- [3] J. COCKLE, *On systems of algebra involving more than one imaginary; and on equations of the fifth degree*, Phil. Mag. (3) 35, 1849, pp. 434–437.
- [4] J. COCKLE, <http://www.oocities.org/cocklebio/>
- [5] YU. A. DROZD, V. V. KIRICHENKO, *Finite dimensional algebras*, Springer, Berlin, 1994, 249 pp.
- [6] I. FRENKEL, M. LIBINE, *Split quaternionic analysis and separation for $SL(2, \mathbb{R})$ and $SL(2, \mathbb{C})/SL(2, \mathbb{R})$* , Advances in Mathematics **228**, (2011), pp 678–763. Also: arXiv:1009.2532v2 [math.RT] 22 Jul 2011, 70 pp.
- [7] K. GÜRLEBECK, W. SPRÖSSIG, *Quaternionic and Clifford calculus for physicists and engineers*, Wiley, Chichester, 1997, 371 pp.
- [8] R. A. HORN, C. R. JOHNSON, *Topics in matrix analysis*, Cambridge University Press, Cambridge, 1991, 607 pp.
- [9] R. A. HORN, C. R. JOHNSON, *Matrix analysis*, Cambridge University Press, Cambridge, 1992, 561 pp.
- [10] D. JANOVSKÁ, G. OPFER, *Linear equations in quaternionic variables*, Mitt. Math. Ges. Hamburg, **27** (2008), pp. 223–234.
- [11] D. JANOVSKÁ, G. OPFER, *Linear equations in quaternions*, Numerical Mathematics and Advanced Applications, Proceedings of ENUMATH 2005, A. B. de Castro, D. Gómez, P. Quintela, and P. Salgado (eds.), Springer, Berlin, Heidelberg, New York, 2006, pp. 945–953, ISBN 3-540-34287-7.
- [12] T. Y. LAM, *The algebraic theory of quadratic forms*, W. A. Benjamin, Reading, Massachusetts, 1973, 343 pp.
- [13] I. NIVEN, *Equations in quaternions*, Amer. Math. Monthly, **48** (1941), pp. 654–661.
- [14] R. S. PIERCE, *Associative algebras*, Springer, New York, 1982, 436 pp.
- [15] M. Sylvester, *Sur l'équations en matrices $px = xq$* , C.R. Acad. Sci. Paris, **99** (1884), pp. 67 – 71, and pp. 115 – 116. [M = Monsieur = J. J.]
- [16] B. L. VAN DER WAERDEN, *Algebra*, 5. Aufl., Springer, Berlin, Göttingen, Heidelberg, 1960, 292 pp.
- [17] F. ZHANG, *Quaternions and matrices of quaternions*, Linear Algebra Appl., **251** (1997), pp. 21–57.